

INVERSE FINITE-TYPE RELATIONS BETWEEN SEQUENCES OF POLYNOMIALS

By

Francisco Marcellán¹ & Ridha Sfaxi²

Abstract

Marcellán, F. & R. Sfaxi: Inverse finite-type relations between sequences of polynomials. Rev. Acad. Colomb. Cienc. **32**(123): 245-255, 2008. ISSN 0370-3908.

Let ϕ be a monic polynomial, with $\deg \phi = t \geq 0$. We say that there is a finite-type relation between two monic polynomial sequences $\{B_n\}_{n \geq 0}$ and $\{Q_n\}_{n \geq 0}$ with respect to ϕ , if there exists $(s, r) \in \mathbb{N}^2$, $r \geq s$, such that

$$\phi(x)Q_n(x) = \sum_{\nu=n-s}^{n+t} \lambda_{n,\nu} B_\nu(x), \quad n \geq s, \quad \text{with } \lambda_{r,r-s} \neq 0. \quad (*)$$

The corresponding inverse finite-type relation of $(*)$ consists in a finite-type relation as follows:

$$\Omega_s^*(x; n)B_n(x) = \sum_{\nu=n-t}^{n+s} \theta_{n,\nu} Q_\nu(x), \quad n \geq t, \quad \text{with } \theta_{r+t,r} \neq 0,$$

where $\deg \Omega_s^*(x; n) = s$, $n \geq t$. When the orthogonality of the two previous sequences is assumed, the inverse finite-type relation is always possible [11]. This work essentially studies the case when only the sequence $\{B_n\}_{n \geq 0}$ is orthogonal. In fact, we find necessary and sufficient conditions leading to inverse finite-type relations. In particular, the structure relation characterizing semi-classical sequences is a special case of the general situation. Some examples will be analyzed.

¹ Departamento de Matemáticas, Universidad Carlos III de Madrid, Avenida de la Universidad 30, 28911 Leganés, Spain. Correo electrónico: pacomarc@ing.uc3m.es

² Département des Méthodes Quantitatives, Institut Supérieur de Gestion de Gabès, Avenue Jilani Habib 6002, Gabès, Tunisie. Correo electrónico: ridhasfaxi@yahoo.fr

Key words: Finite-type relations, recurrence relations, orthogonal polynomials, semi-classical polynomials.

Resumen

Sea ϕ un polinomio mónico, con $\deg \phi = t \geq 0$. Decimos que hay relación de tipo finito entre dos sucesiones de polinomios mónicos $\{B_n\}_{n \geq 0}$ y $\{Q_n\}_{n \geq 0}$ con respecto a ϕ , si existe $(s, r) \in \mathbb{N}^2$, $r \geq s$, tal que

$$\phi(x)Q_n(x) = \sum_{\nu=n-s}^{n+t} \lambda_{n,\nu} B_\nu(x), \quad n \geq s, \text{ with } \lambda_{r,r-s} \neq 0. \quad (*)$$

La correspondiente relación de tipo finito de (*) consiste en una relación de tipo finito como sigue:

$$\Omega_s^*(x; n)B_n(x) = \sum_{\nu=n-t}^{n+s} \theta_{n,\nu}^* Q_\nu(x), \quad n \geq t, \text{ with } \theta_{r+t,r}^* \neq 0,$$

donde $\deg \Omega_s^*(x; n) = s$, $n \geq t$. Cuando se supone la ortogonalidad de las dos sucesiones previas, la relación de tipo finito inversa siempre es posible [11]. En este trabajo se estudia el caso en que solo la sucesión $\{B_n\}_{n \geq 0}$ es ortogonal. De hecho, encontramos condiciones necesarias y suficientes que conducen a relaciones de tipo finito inversas. En particular, la la relación de estructura que caracteriza a las sucesiones semiclásicas es un caso especial de la situación general. Se estudian varios ejemplos.

Palabras clave: Relaciones de tipo finito, relaciones de recurrencia, polinomios ortogonales, polinomios semi clásicos.

1. Introduction and background

Let \mathbb{P} be the linear space of complex polynomials in one variable and \mathbb{P}' its topological dual space. We denote by $\langle u, f \rangle$ the action of $u \in \mathbb{P}'$ on $f \in \mathbb{P}$ and by $(u)_n := \langle u, x^n \rangle$, $n \geq 0$, the moments of u with respect to the polynomial sequence $\{x^n\}_{n \geq 0}$.

We will introduce some useful operations in \mathbb{P}' . For any linear functional u and any polynomial h , let $Du = u'$ and hu be the linear functionals defined by duality

$$\begin{aligned} \langle u', f \rangle &:= -\langle u, f' \rangle, & f &\in \mathbb{P}, \\ \langle hu, f \rangle &:= \langle u, hf \rangle, & f, h &\in \mathbb{P}. \end{aligned}$$

Let $\{B_n\}_{n \geq 0}$ be a monic polynomial sequence (MPS), $\deg B_n = n$, $n \geq 0$, and $\{u_n\}_{n \geq 0}$ its dual sequence, $u_n \in \mathbb{P}'$, $n \geq 0$, defined by $\langle u_n, B_m \rangle := \delta_{n,m}$, $n, m \geq 0$, where $\delta_{n,m}$ is the Kronecker symbol.

Let recall the following results [11].

Lemma 1.1. For any $u \in \mathbb{P}'$ and any integer $m \geq 1$, the following statements are equivalent.

$$i) \langle u, B_{m-1} \rangle \neq 0, \quad \langle u, B_n \rangle = 0, \quad n \geq m.$$

$$ii) \text{ There exist } \lambda_\nu \in \mathbb{C}, \quad 0 \leq \nu \leq m-1, \quad \lambda_{m-1} \neq 0, \\ \text{ such that } u = \sum_{\nu=0}^{m-1} \lambda_\nu u_\nu.$$

As a consequence, the dual sequence $\{u_n^{[1]}\}_{n \geq 0}$ of the sequence $\{B_n^{[1]}\}_{n \geq 0}$, where $B_n^{[1]}(x) = (n+1)^{-1}B'_{n+1}(x)$, $n \geq 0$, satisfies

$$(u_n^{[1]})' = -(n+1)u_{n+1}, \quad n \geq 0. \quad (1.1)$$

Definition 1.2. The linear functional u is said to be regular if there exists a monic polynomial sequence $\{B_n\}_{n \geq 0}$ such that

$$\langle u, B_n B_m \rangle = b_n \delta_{n,m}, \quad n, m \geq 0, \quad (1.2)$$

where

$$b_n = \langle u, B_n^2 \rangle \neq 0, \quad n \geq 0. \quad (1.3)$$

Then the sequence $\{B_n\}_{n \geq 0}$ is said to be orthogonal (MOPS) with respect to u .

As a straightforward consequence we get

• The linear functional can be represented by $u = (u)_0 u_0$, and the following relations hold

$$u_n = b_n^{-1} B_n u, \quad n \geq 0. \quad (1.4)$$

•• The sequence $\{B_n\}_{n \geq 0}$ satisfies the three-term recurrence relation

$$\begin{aligned} B_{n+2}(x) &= (x - \beta_{n+1})B_{n+1}(x) - \gamma_{n+1}B_n(x), \quad n \geq 0, \\ B_1(x) &= x - \beta_0, \quad B_0(x) = 1, \end{aligned} \tag{1.5}$$

where $\gamma_{n+1} \neq 0, n \geq 0$ (see [4]).

In the sequel and under the assumption of the previous definition, we need to put

$$b_{n,m}^\nu = b_m^{-1} \langle u, x^\nu B_m B_n \rangle, \quad (n, \nu, m) \in \mathbb{N}^3. \tag{1.6}$$

In particular, one has

$$b_{n,m}^\nu = \begin{cases} 0, & \text{if } \nu + m < n, \quad 0 \leq m < n, \nu \geq 0, \\ (b_n/b_m), & \text{if } \nu = n - m, \quad 0 \leq m \leq n. \end{cases}$$

Let ϕ be a monic polynomial, with $\deg \phi = t \geq 0$. For any MPS $\{B_n\}_{n \geq 0}$ and $\{Q_n\}_{n \geq 0}$ with dual sequences $\{u_n\}_{n \geq 0}$ and $\{v_n\}_{n \geq 0}$ respectively, the following formula always holds

$$\phi(x)Q_n(x) = \sum_{\nu=0}^{n+t} \lambda_{n,\nu} B_\nu(x), \quad n \geq 0, \tag{1.7}$$

where $\lambda_{n,\nu} = \langle u_\nu, \phi Q_n \rangle, 0 \leq \nu \leq n+t, n \geq 0$.

Definition 1.3. ([12]) If there exists an integer $s \geq 0$ such that

$$\phi(x)Q_n(x) = \sum_{\nu=n-s}^{n+t} \lambda_{n,\nu} B_\nu(x), \quad n \geq s, \tag{1.8}$$

and

$$\exists r \geq s, \lambda_{r,r-s} \neq 0, \tag{1.9}$$

then, we shall say that (1.8) – (1.9) gives a finite-type relation between $\{B_n\}_{n \geq 0}$ and $\{Q_n\}_{n \geq 0}$, with respect to ϕ .

When instead of (1.9), we take

$$\lambda_{n,n-s} \neq 0, \quad n \geq s, \tag{1.9'}$$

we shall say that (1.8) – (1.9') is a strictly finite-type relation.

The corresponding inverse finite-type relation of (1.8) – (1.9) consists in establishing, whenever it is possible, a finite-type relation between $\{Q_n\}_{n \geq 0}$ and $\{B_n\}_{n \geq 0}$, as follows

$$\Omega_s^*(x; n)B_n(x) = \sum_{\nu=n-t}^{n+s} \theta_{n,\nu}^* Q_\nu(x), \quad n \geq t, \tag{1.10}$$

$\theta_{r+t,r}^* \neq 0$, where $\{\Omega_s^*(x; n)\}_{n \geq t}$ is a MPS, $\deg \Omega_s^*(x; n) = s, n \geq t$, and

$$(\theta_{n,\nu}^*)_{\nu=n-t}^{n+s}, \quad n \geq t, \tag{1.11}$$

a system of complex numbers (SCN), with $\theta_{n,n+s}^* = 1, n \geq t$.

When both two sequences are orthogonal, the inverse relation is always possible. In this case, the polynomials $\Omega_s^*(x; n), n \geq 0$, are independent of n , (see [12], Proposition 2.4). As a current example, we can mention the two structure relations characterizing the classical polynomials, (**Hermite, Laguerre, Bessel, Jacobi**, see [11]), which could solely be two inverse finite-type relations.

In other studies, we find several situations where one of the two sequences is orthogonal. For example, the structure relations characterizing semi-classical sequences associated with Hahn's operators $L_{q,\omega}$, with parameters q and ω , [9]. The Coherent pairs and Diagonal sequences are also examples of finite type-relations [7, 12, 13, 14]. But the inverse relations corresponding to other finite-type relations are not yet considered.

The paper essentially gives a necessary and sufficient condition allowing the existence of the inverse finite-type relations when the orthogonality of the sequence $\{B_n\}_{n \geq 0}$ is assumed. From now on, it would be necessary to study the case where the sequence $\{Q_n\}_{n \geq 0}$ is orthogonal. It would be very useful to deal with many other situations like General Coherent pairs, see [6, 8] in the framework of Sobolev inner products.

2. A basic result

We use this section to introduce some auxiliary result for the proof of the main theorem in section 3.

Lemma 2.1. Suppose $\{B_n\}_{n \geq 0}$ is a MOPS and $\{Q_n\}_{n \geq 0}$ fulfils (1.8) – (1.9), where $t = 0$ and $s \geq 1$. For any SCN $(\theta_{n,\nu}^*)_{\nu=n}^{n+s}, n \geq 0$, where $\theta_{n,n+s} = 1, n \geq 0$, and $\theta_{r,r} \neq 0$, there exist a unique MPS $\{\Omega_s(x; n)\}_{n \geq 0}$, $\deg \Omega_s(x; n) = s, n \geq 0$, and a SCN $(\zeta_{n,\nu}^{[0]})_{\nu=n-s}^{n-1}, n \geq 0$, such that

$$\begin{aligned} \sum_{\nu=n}^{n+s} \theta_{n,\nu}^* Q_\nu(x) &= \sum_{i=n-s}^{n+s} \theta_{n,i}^{[0]} B_i(x) \\ &= \Omega_s(x; n)B_n(x) + \sum_{\nu=n-s}^{n-1} \zeta_{n,\nu}^{[0]} B_\nu(x), \quad n \geq 0, \end{aligned} \tag{2.1}$$

where

$$\theta_{n,i}^{[0]} = \sum_{\nu=\max(n,i)}^{\min(n,i)+s} \theta_{n,\nu} \lambda_{\nu,i}, \quad n-s \leq i \leq n+s, n \geq 0, \quad (2.2)$$

$$\theta_{r,r-s}^{[0]} = \theta_{r,r} \lambda_{r,r-s} \neq 0, \quad (2.3)$$

$$\sum_{\nu=n}^{m+s} \theta_{n,\nu} \lambda_{\nu,m} = b_m^{-1} \langle u, \Omega_s(x; n) B_n B_m \rangle + \zeta_{n,m}^{[0]}, \quad n-s \leq m \leq n-1, n \geq 0, \quad (2.4)$$

$$\sum_{\nu=m}^{n+s} \theta_{n,\nu} \lambda_{\nu,m} = b_m^{-1} \langle u, \Omega_s(x; n) B_n B_m \rangle, \quad n \leq m \leq n+s-1, n \geq 0. \quad (2.5)$$

Proof. Let $(\theta_{n,\nu})_{\nu=n}^{n+s}, n \geq 0$, where $\theta_{n,n+s} = 1, n \geq 0$, and $\theta_{r,r} \neq 0$, be a SCN. From (1.8) – (1.9), with $t = 0$ and $s \geq 1$, we get

$$\begin{aligned} \sum_{\nu=n}^{n+s} \theta_{n,\nu} Q_{\nu}(x) &= \sum_{\nu=n}^{n+s} \theta_{n,\nu} \sum_{i=\nu-s}^{\nu} \lambda_{\nu,i} B_i(x) \\ &= \sum_{\nu=n}^{n+s} \theta_{n,\nu} \sum_{i=n-s}^{n+s} \chi_{i,\nu} \lambda_{\nu,i} B_i(x), \quad n \geq 0, \end{aligned}$$

where, for each pair of integers (i, ν) such that $n-s \leq i \leq n+s$ and $n \leq \nu \leq n+s$, we took

$$\chi_{i,\nu} = \begin{cases} 1, & \text{if } \nu-s \leq i \leq \nu, \\ 0, & \text{otherwise.} \end{cases}$$

The permutation of these two sums yields

$$\sum_{\nu=n}^{n+s} \theta_{n,\nu} Q_{\nu}(x) = \sum_{i=n-s}^{n+s} \theta_{n,i}^{[0]} B_i(x),$$

where

$$\theta_{n,i}^{[0]} = \sum_{\nu=\max(n,i)}^{\min(i+s,n+s)} \theta_{n,\nu} \lambda_{\nu,i}, \quad n-s \leq i \leq n+s, n \geq 0, \quad (2.6)$$

$$\theta_{r,r-s}^{[0]} = \theta_{r,r} \lambda_{r,r-s} \neq 0.$$

Hence, (2.2) and (2.3) are valid.

The Euclidean division by $B_n(x)$ in the right hand side in (2.6) gives

$$\sum_{i=n-s}^{n+s} \theta_{n,i}^{[0]} B_i(x) = \Omega_s(x; n) B_n(x) + \sum_{\nu=0}^{n-1} \zeta_{n,\nu}^{[0]} B_{\nu}(x), \quad n \geq 0.$$

Multiplying by $B_m(x)$ and using the orthogonality of $\{B_n\}_{n \geq 0}$,

$$\sum_{i=n-s}^{n+s} \theta_{n,i}^{[0]} \delta_{m,i} = b_m^{-1} \langle u, \Omega_s(x; n) B_n B_m \rangle + \sum_{\nu=0}^{n-1} \zeta_{n,\nu}^{[0]} \delta_{m,\nu}.$$

In particular, for $0 \leq m \leq n-s-1$ and $n \geq s+1$, it follows that $\zeta_{n,m}^{[0]} = 0$. Hence, (2.1) holds. Moreover, for $n-s \leq m \leq n-1$ and $n \geq s$, we recover (2.4).

Finally, for $n \leq m \leq n+s-1$ and $n \geq 0$, we deduce (2.5). \square

Proposition 2.2. Assume $\{B_n\}_{n \geq 0}$ is a MOPS and $\{Q_n\}_{n \geq 0}$ fulfils (1.8) – (1.9), with $t \geq 1$. For any SCN $(\theta_{n,\nu})_{\nu=n-t}^{n+s}, n \geq 0$, where $\theta_{n,n+s} = 1, n \geq 0$ and $\theta_{r+t,r} \neq 0$, there exist a unique MPS $\{\Omega_{s+t}(x; n)\}_{n \geq 0}$, where $\deg \Omega_{s+t}(x; n) = s+t, n \geq 0$, and a SCN $(\zeta_{n,\nu}^{[t]})_{\nu=n-s-t}^{n-1}, n \geq 0$, such that for every integer $n \geq 0$

$$\begin{aligned} \phi(x) \sum_{\nu=n-t}^{n+s} \theta_{n,\nu} Q_{\nu}(x) &= \sum_{i=n-s-t}^{n+s+t} \theta_{n,i}^{[t]} B_i(x) \\ &= \Omega_{s+t}(x; n) B_n(x) \\ &\quad + \sum_{\nu=n-s-t}^{n-1} \zeta_{n,\nu}^{[t]} B_{\nu}(x), \quad (2.7) \end{aligned}$$

where

$$\theta_{n,i}^{[t]} = \sum_{\nu=\max(n,i)}^{\min(n,i)+s+t} \theta_{n,\nu-t} \lambda_{\nu-t,i}, \quad n-s-t \leq i \leq n+s+t, \quad (2.8)$$

$$\theta_{r+t,r-s}^{[t]} = \theta_{r+t,r} \lambda_{r,r-s} \neq 0, \quad (2.9)$$

$$\sum_{\nu=n}^{m+s+t} \theta_{n,\nu-t} \lambda_{\nu-t,m} = b_m^{-1} \langle u, \Omega_{s+t}(x; n) B_n B_m \rangle + \zeta_{n,m}^{[t]}, \quad n-s-t \leq m \leq n-1, \quad (2.10)$$

$$\sum_{\nu=m}^{n+s+t} \theta_{n,\nu-t} \lambda_{\nu-t,m} = b_m^{-1} \langle u, \Omega_{s+t}(x; n) B_n B_m \rangle, \quad n \leq m \leq n+s+t-1. \quad (2.11)$$

Proof. The case $t = 0$ was analyzed in Lemma 2.1. Let us take $t \geq 1$. Consider the MPS $\{P_n\}_{n \geq t}$ defined by

$$P_{n+t}(x) = \phi(x) Q_n(x), \quad n \geq 0, \quad (2.12)$$

From (1.8) – (1.9), we have

$$P_n(x) = \sum_{\nu=n-t-s}^n \tilde{\lambda}_{n,\nu} B_\nu(x), \quad n \geq t,$$

where $\tilde{\lambda}_{n,\nu} = \lambda_{n-t,\nu}$, $n-t-s \leq \nu \leq n$, $n \geq t$, and $\tilde{\lambda}_{t+r,r-s} \neq 0$. Now, let $(\theta_{n,\nu})_{\nu=n-t}^{n+s}$, $n \geq 0$, where $\theta_{n,n+s} = 1$, $n \geq 0$, and $\theta_{r+t,r} \neq 0$, be a SCN. One has

$$\begin{aligned} \phi(x) \sum_{\nu=n-t}^{n+s} \theta_{n,\nu} Q_\nu(x) &= \sum_{\nu=n-t}^{n+s} \theta_{n,\nu} P_{\nu+t}(x) \\ &= \sum_{\nu=n}^{n+t+s} \tilde{\theta}_{n,\nu} P_\nu(x), \quad n \geq 0, \end{aligned} \quad (2.13)$$

where $\tilde{\theta}_{n,\nu} = \theta_{n,\nu-t}$, $n \leq \nu \leq n+t+s$, $n \geq 0$. Obviously, $(\tilde{\theta}_{n,\nu})_{\nu=n}^{n+t+s}$, $n \geq 0$, is a SCN such that

$$\tilde{\theta}_{n,n+t+s} = \theta_{n,n+s} = 1, \quad n \geq 0, \quad \tilde{\theta}_{r+t,r+t} = \theta_{r+t,r} \neq 0.$$

But from Lemma 2.1, there exist a unique MPS $\{\Omega_{t+s}(x;n)\}_{n \geq 0}$ and a SCN $(\zeta_{n,\nu}^{[t]})_{\nu=n-t-s}^{n-1}$, $n \geq 0$, such that

$$\begin{aligned} \sum_{\nu=n}^{n+t+s} \tilde{\theta}_{n,\nu} P_\nu(x) &= \sum_{i=n-t-s}^{n+t+s} \theta_{n,i}^{[t]} B_i(x) \\ &= \Omega_{t+s}(x;n) B_n(x) + \sum_{\nu=n-t-s}^{n-1} \zeta_{n,\nu}^{[t]} B_\nu(x), \end{aligned} \quad (2.14)$$

for every integer $n \geq 0$, where

$$\theta_{n,i}^{[t]} = \sum_{\nu=\max(n,i)}^{\min(n,i)+t+s} \tilde{\theta}_{n,\nu} \tilde{\lambda}_{\nu,i}, \quad n-t-s \leq i \leq n+t+s,$$

$$\theta_{r+t,r-s}^{[t]} = \tilde{\theta}_{r+t,r+t} \tilde{\lambda}_{r+t,r-s} \neq 0,$$

$$\begin{aligned} \sum_{\nu=n}^{m+t+s} \tilde{\theta}_{n,\nu} \tilde{\lambda}_{\nu,m} &= b_m^{-1} \langle u, \Omega_{t+s}(x;n) B_n B_m \rangle + \zeta_{n,m}^{[t]}, \\ n-t-s &\leq m \leq n-1, \end{aligned}$$

$$\begin{aligned} \sum_{\nu=m}^{n+t+s} \tilde{\theta}_{n,\nu} \tilde{\lambda}_{\nu,m} &= b_m^{-1} \langle u, \Omega_{t+s}(x;n) B_n B_m \rangle, \\ n &\leq m \leq n+t+s-1. \end{aligned}$$

Finally, by using (2.13), (2.14), and taking into account the expressions of $\tilde{\lambda}_{n,\nu}$ and $\tilde{\theta}_{n,\nu}$, we find the desired results. \square

3. A matrix approach and main results

In this section, we will work under the assumptions of the Proposition 2.2 and we will give a matrix approach to our problem.

If $\Omega_{t+s}(x;n) = \sum_{\nu=0}^{t+s} v_{n,\nu} x^\nu$, $n \geq 0$, where $v_{n,t+s} = 1$, then relation (2.10) reads

$$\sum_{\nu=n}^{m+s+t} \lambda_{\nu-t,m} \theta_{n,\nu-t} = \sum_{\nu=0}^{t+s-1} b_{n,m}^\nu v_{n,\nu} + \zeta_{n,m}^{[t]} + b_{n,m}^{s+t}, \quad n-s-t \leq m \leq n-1,$$

or, alternatively,

$$\begin{aligned} \sum_{j=1}^{m+s+t-n+1} \lambda_{j+n-t-1,m} \theta_{n,j+n-t-1} \\ = \sum_{j=1}^{t+s} b_{n,m}^{j-1} v_{n,j-1} + \zeta_{n,m}^{[t]} + b_{n,m}^{s+t}, \end{aligned}$$

for every $n-s-t \leq m \leq n-1$.

Replacing m by $i+n-s-t-1$, we get

$$\begin{aligned} \sum_{j=1}^i k_{i,j}^n \Theta_{n,j} = \sum_{j=1}^{t+s} t_{i,j}^n V_{n,j} + \zeta_{n,i+n-s-t-1}^{[t]} \\ b_{n,i+n-s-t-1}^{s+t}, \quad 1 \leq i \leq s+t, \end{aligned}$$

where for $i, j = 1, 2, \dots, s+t$,

$$k_{i,j}^n = \begin{cases} \lambda_{j+n-t-1,i+n-s-t-1}, & 1 \leq j \leq i \\ 0, & \text{otherwise,} \end{cases}$$

and $t_{i,j}^n = b_{n,i+n-s-t-1}^{j-1}$,

$$\Theta_{n,j} = \theta_{n,j+n-t-1}, \quad \text{and} \quad V_{n,j} = v_{n,j-1}.$$

So we can write it as

$$\mathbf{K}_n \Theta_n = \mathbf{T}_n V_n + W_n + E_n, \quad n \geq 0, \quad (3.1)$$

where

$$\mathbf{K}_n = (k_{i,j}^n)_{1 \leq i,j \leq s+t}, \quad \mathbf{T}_n = (t_{i,j}^n)_{1 \leq i,j \leq s+t},$$

$$\Theta_n = (\Theta_{n,1}, \Theta_{n,2}, \dots, \Theta_{n,s+t})^T,$$

$$V_n = (V_{n,1}, V_{n,2}, \dots, V_{n,s+t})^T,$$

$$W_n = (\zeta_{n,n-s-t}^{[t]}, \zeta_{n,n+1-s-t}^{[t]}, \dots, \zeta_{n,n-1}^{[t]})^T, \quad \text{and}$$

$$E_n = (b_{n,n-s-t}^{s+t}, b_{n,n+1-s-t}^{s+t}, \dots, b_{n,n-1}^{s+t})^T.$$

In the same way, using $\theta_{n,n+s} = 1$, (2.11) can be written as

$$\sum_{\nu=m}^{n+s+t-1} \lambda_{\nu-t,m} \theta_{n,\nu-t} = \sum_{\nu=0}^{t+s-1} b_{n,m}^{\nu} v_{n,\nu} + b_{n,m}^{s+t} - \lambda_{n+s,m},$$

$$n \leq m \leq n + s + t - 1,$$

or, equivalently,

$$\sum_{j=m-n+1}^{s+t} \lambda_{j+n-t-1,m} \theta_{n,j+n-t-1} =$$

$$\sum_{j=1}^{t+s} b_{n,m}^{j-1} v_{n,j-1} + b_{n,m}^{s+t} - \lambda_{n+s,m},$$

for every $n \leq m \leq n+s+t-1$. Replacing m by $i+n-1$, we get

$$\sum_{j=i-1}^{t+s} m_{i,j}^n \Theta_{n,j} = \sum_{j=1}^{t+s} s_{i,j}^n V_{n,j} + b_{n,i+n-1}^{s+t} - \lambda_{n+s,i+n-1},$$

$$1 \leq i \leq s+t,$$

where for $i, j = 1, 2, \dots, s+t$,

$$m_{i,j}^n := \begin{cases} \lambda_{j+n-t-1,i+n-1}, & 1 \leq i \leq j \\ 0, & \text{otherwise,} \end{cases}$$

$$s_{i,j}^n := \begin{cases} b_{n,i+n-1}^{j-1}, & 1 \leq i \leq j \\ 0, & \text{otherwise.} \end{cases}$$

Thus, we can use the matrix representation

$$M_n \Theta_n = S_n V_n + F_n, \quad n \geq 0, \quad (3.2)$$

where

$$M_n = (m_{i,j}^n)_{1 \leq i,j \leq s+t}, \quad S_n = (s_{i,j}^n)_{1 \leq i,j \leq s+t}$$

and

$$F_n = (b_{n,n}^{s+t} - \lambda_{n+s,n}, b_{n,n+1}^{s+t} - \lambda_{n+s,n+1}, \dots, b_{n,n+s+t-1}^{s+t} - \lambda_{n+s,n+s+t-1})^T.$$

Our data are $\Theta_n, E_n, F_n, M_n, S_n, T_n, K_n$ and our unknowns are V_n and W_n .

From (3.2), we get

$$V_n = S_n^{-1}(M_n \Theta_n - F_n). \quad (3.3)$$

Thus, substituting in (3.1) we get $K_n \Theta_n - W_n - E_n = T_n S_n^{-1}(M_n \Theta_n - F_n)$, i.e,

$$W_n = (K_n - T_n S_n^{-1} M_n) \Theta_n + T_n S_n^{-1} F_n - E_n.$$

As a consequence, for every choice of Θ_n , we get W_n . From (3.3), we deduce V_n .

On the other hand, there exists a one-to-one correspondence between the vectors W_n and Θ_n if and only if the matrix of dimension $s+t$, $K_n - T_n S_n^{-1} M_n$, is nonsingular.

Under such a condition, there exists a unique choice for Θ_n such that $W_n = 0$. Thus, we get

$$\Theta_n = (K_n - T_n S_n^{-1} M_n)^{-1} (E_n - T_n S_n^{-1} F_n),$$

and from (3.3), $V_n = S_n^{-1} M_n \Theta_n - S_n^{-1} F_n$. Then,

$$V_n = (K_n M_n^{-1} S_n - T_n)^{-1} E_n - [(K_n M_n^{-1} S_n - T_n)^{-1} T_n + I_{s+t}] S_n^{-1} F_n,$$

where I_{s+t} is the unit matrix. Hence, the polynomial $\Omega_{s+t}(x; n)$ is explicitly given.

Let introduce

$$\Delta_n(t, s) = \det(K_n - T_n S_n^{-1} M_n), \quad n \geq 0.$$

Thus, we have proved the following result

Proposition 3.1. Assume $\{B_n\}_{n \geq 0}$ is a MOPS and $\{Q_n\}_{n \geq 0}$ fulfils (1.8)–(1.9). For a fixed integer $p \geq t+1$, the following statements are equivalent.

- i) $\Delta_n(t, s) \neq 0, n \geq p$.
- ii) There exist a unique SCN $(\theta_{n,\nu}^*)_{\nu=n-t}^{n+s}, n \geq p$, with $\theta_{n,n+s}^* = 1, n \geq p$, and $\theta_{r+t,r}^* \neq 0$, if $p \leq r+t$, and a unique MPS $\{\Omega_{s+t}^*(x; n)\}_{n \geq p}$, $\deg \Omega_{s+t}^*(x; n) = s+t, n \geq p$, such that

$$\Omega_{s+t}^*(x; n) B_n(x) = \phi(x) \sum_{\nu=n-t}^{n+s} \theta_{n,\nu}^* Q_\nu(x), \quad (3.4)$$

for $n \geq p$.

Our main result is

Theorem 3.2. Let $\{B_n\}_{n \geq 0}$ be a MOPS and $\{Q_n\}_{n \geq 0}$ be the MPS satisfying (1.8)–(1.9). For each fixed integer $p \geq t+1$, if we suppose that $\phi(x)$ and $B_n(x)$ are coprime for every $n \geq p$, then the following statements are equivalent.

- i) $\Delta_n(t, s) \neq 0, n \geq p$.
- ii) There exist a unique SCN $(\theta_{n,\nu}^*)_{\nu=n-t}^{n+s}, n \geq p$, where $\theta_{n,n+s}^* = 1, n \geq p$, and $\theta_{r+t,r}^* \neq 0$ if $p \leq r+t$, and a unique MPS $\{\Omega_s^*(x; n)\}_{n \geq p}$, $\deg \Omega_s^*(x; n) = s, n \geq p$, such that

$$\Omega_s^*(x; n) B_n(x) = \sum_{\nu=n-t}^{n+s} \theta_{n,\nu}^* Q_\nu(x), \quad n \geq p. \quad (3.5)$$

Proof. Taking into account $\phi(x)$ and $B_n(x)$ are coprime for every $n \geq p$, from (3.4) we deduce that ϕ divides $\Omega_{s+t}^*(x; n)$, $n \geq p$. So, $\Omega_{s+t}^*(x; n) = \phi(x)\Omega_s^*(x; n)$, $n \geq p$. Hence, the desired result follows. \square

The orthogonal polynomial sequence $\{B_n\}_{n \geq 0}$ and the polynomial sequence $\{Q_n\}_{n \geq 0}$ can be related by a general finite-type relation (see [1]). It reads as follows

$$F(Q_n, \dots, Q_{n-l}) = G(B_n, \dots, B_{n-s}),$$

where F and G are fixed functions.

When F and G are linear functions, some situations dealing with the inverse problem have been analyzed in [1,2]. There, necessary and sufficient conditions in order to $\{Q_n\}_{n \geq 0}$ be orthogonal are obtained.

This kind of linear relations reads as follows.

There exists $(l, s, r) \in \mathbb{N}^3$, with $r \geq \tilde{s} = \max(l, s)$ such that

$$\sum_{\nu=n-l}^n \zeta_{n,\nu} Q_\nu(x) = \sum_{\nu=n-s}^n \lambda_{n,\nu} B_\nu(x), \quad n \geq \tilde{s}, \quad (3.6)$$

with $\zeta_{r,r-l} \lambda_{r,r-s} \neq 0$. Here, $\zeta_{n,n} = \lambda_{n,n} = 1$, $n \geq \tilde{s}$.

More recently, in [5], **A. M. Delgado** and **F. Marcellán** exhaustively describe all the set of pairs of quasi-definite (regular) linear functionals such that their corresponding sequences of monic polynomials $\{P_n\}_{n \geq 0}$ and $\{R_n\}_{n \geq 0}$ are related by a differential expression

$$P_n(x) + s_n P_{n-1}(x) = R_n^{[1]}(x) + t_n R_{n-1}^{[1]}(x), \quad n \geq 1,$$

where $t_n \neq 0$, for every $n \geq 1$, and with the technical condition $t_1 \neq s_1$.

Notice that in general $\{R_n^{[1]}\}_{n \geq 0}$ is not a MOPS.

In the same context of our contribution, we show that the corresponding inverse finite-type relation between two sequences satisfying (3.6) is possible under certain conditions.

Indeed, let consider the MPS $\{C_n\}_{n \geq \tilde{s}}$ given by

$$C_n(x) = \sum_{\nu=n-s}^n \lambda_{n,\nu} B_\nu(x), \quad n \geq \tilde{s}. \quad (3.7)$$

With the finite-type relation between the sequences $\{C_n\}_{n \geq \tilde{s}}$ and $\{B_n\}_{n \geq \tilde{s}}$, we can associate the determinants $\Delta_n(0, s)$, $n \geq \tilde{s}$. So, we have.

Corollary 3.3. *Let $\{B_n\}_{n \geq 0}$ be a MOPS and $\{Q_n\}_{n \geq 0}$ be the MPS satisfying (3.6). For each fixed integer*

$p \geq \max(s, l, 1)$, if $\Delta_n(0, s) \neq 0$, $n \geq p$, then there exist a unique SCN $(\zeta_{n,\nu}^*)_{\nu=n-l}^{n+s}$, $n \geq p$, where $\zeta_{n,n+s}^* = 1$, $n \geq p$, and $\zeta_{r,r-l}^* \neq 0$ if $p \leq r$, and a unique MPS $\{\Omega_s^*(x; n)\}_{n \geq p}$, $\deg \Omega_s^*(x; n) = s$, $n \geq p$, such that

$$\Omega_s^*(x; n) B_n(x) = \sum_{\nu=n-l}^{n+s} \zeta_{n,\nu}^* Q_\nu(x), \quad n \geq p. \quad (3.8)$$

Proof. From Theorem 3.2, with $t = 0$, there exists the corresponding inverse finite-type relation associated with the relation (3.7) if and only if $\Delta_n(0, s) \neq 0$, $n \geq p$. Equivalently, there exist a unique SCN $(\theta_{n,\nu}^*)_{\nu=n}^{n+s}$, $n \geq p$, where $\theta_{n,n+s}^* = 1$, $n \geq p$, and $\theta_{r,r}^* \neq 0$, if $p \leq r$, and a unique MPS $\{\Omega_s^*(x; n)\}_{n \geq p}$, $\deg \Omega_s^*(x; n) = s$, $n \geq p$, such that

$$\Omega_s^*(x; n) B_n(x) = \sum_{\nu=n}^{n+s} \theta_{n,\nu}^* C_\nu(x), \quad n \geq p. \quad (3.9)$$

But from (3.6) and (3.7), the above expression becomes

$$\begin{aligned} \Omega_s^*(x; n) B_n(x) &= \sum_{\nu=n}^{n+s} \theta_{n,\nu}^* \sum_{i=\nu-l}^{\nu} \zeta_{\nu,i} Q_i(x) \\ &= \sum_{\nu=n}^{n+s} \theta_{n,\nu}^* \sum_{i=n-l}^{n+s} \tilde{\chi}_{i,\nu} \zeta_{\nu,i} Q_i(x), \quad n \geq p, \end{aligned}$$

where, for each pair of integers (i, ν) such that $n-l \leq i \leq \nu$ and $n \leq \nu \leq n+s$, we took

$$\tilde{\chi}_{i,\nu} = \begin{cases} 1, & \text{if } \nu-l \leq i \leq \nu, \\ 0, & \text{otherwise.} \end{cases}$$

The permutation inside these two sums yields

$$\Omega_s^*(x; n) B_n(x) = \sum_{i=n-l}^{n+s} \zeta_{n,i}^* Q_i(x),$$

where

$$\zeta_{n,i}^* = \sum_{\nu=\max(n,i)}^{\min(i+l,n+s)} \theta_{n,\nu}^* \zeta_{\nu,i}$$

if $n-l \leq i \leq n+s$, $n \geq p$, and

$$\zeta_{r,r-l}^* = \theta_{r,r}^* \zeta_{r,r-l} \neq 0,$$

if $p \leq r$. \square

4. The case: $(t, s) = (0, 1)$

Let $\{B_n\}_{n \geq 0}$ be a MOPS with respect to the linear functional u and satisfying the three-term recurrence relation (1.5).

Consider the following finite-type relation between $\{B_n\}_{n \geq 0}$ and $\{Q_n\}_{n \geq 0}$, with index $s = 1$, with respect to $\phi(x) = 1$,

$$Q_n(x) = B_n(x) + \lambda_{n,n-1}B_{n-1}(x), \quad n \geq 0, \quad (4.1)$$

$$\exists r \geq 1, \quad \lambda_{r,r-1} \neq 0. \quad (4.2)$$

From Lemma 2.1, for every set of complex numbers, $\theta_{n,n}$, $n \geq 0$, with $\theta_{r,r} \neq 0$, there exists a unique MPS $\{\Omega_1(x; n)\}_{n \geq 0}$, where $\Omega_1(x; n) = x + v_{n,0}$, $n \geq 0$, and a unique set of complex numbers, $\zeta_{n,n-1}^{[0]}$, $n \geq 0$, such that

$$Q_{n+1}(x) + \theta_{n,n}Q_n(x) = \Omega_1(x; n)B_n(x) + \zeta_{n,n-1}^{[0]}B_{n-1}(x), \quad n \geq 0, \quad (4.3)$$

where

$$\begin{cases} \lambda_{n,n-1}\theta_{n,n} &= \zeta_{n,n-1}^{[0]} + \gamma_n, \quad n \geq 1, \\ \theta_{n,n} - v_{n,0} &= -\lambda_{n+1,n} + \beta_n, \quad n \geq 0. \end{cases} \quad (4.4)$$

The determinants associated with (4.1) – (4.2) are given by

$$\Delta_0(0, 1) = 0, \quad \Delta_n(0, 1) = \lambda_{n,n-1}, \quad n \geq 1, \quad (4.5)$$

where $\Delta_r(0, 1) = \lambda_{r,r-1} \neq 0$. As a consequence of Theorem 3.2, when $t = 0$ and $s = 1$, we have the following result

Proposition 4.1. *Let $\{B_n\}_{n \geq 0}$ be a MOPS and $\{Q_n\}_{n \geq 0}$ be the MPS satisfying (4.1) – (4.2). For every fixed integer $p \geq 1$, the following statements are equivalent*

- i) $\lambda_{n,n-1} \neq 0$, $n \geq p$.
- ii) There exist a unique set of complex numbers $\theta_{n,n}^* \neq 0$, $n \geq p$, and a unique MPS $\{\Omega_1^*(x; n)\}_{n \geq p}$, $\deg \Omega_1^*(x; n) = 1$, $n \geq p$, such that

$$\Omega_1^*(x; n)B_n(x) = Q_{n+1}(x) + \theta_{n,n}^*Q_n(x), \quad n \geq p. \quad (4.6)$$

We write

$$\theta_{n,n}^* = \frac{\gamma_n}{\lambda_{n,n-1}}, \quad n \geq p, \quad (4.7)$$

$$\Omega_1^*(x; n) = x + v_{n,0}^*,$$

where

$$v_{n,0}^* = \frac{\gamma_n}{\lambda_{n,n-1}} + \lambda_{n+1,n} - \beta_n, \quad n \geq p. \quad (4.8)$$

Example. In order to illustrate the result of Proposition 4.1, we study the structure relation characterizing a semi-classical polynomial sequence, $\{B_n\}_{n \geq 0}$, orthogonal with respect to the linear functional u solution of the functional equation

$$u' + \psi u = 0, \quad (4.9)$$

where $\psi(x) = -ix^2 + 2x - i(\alpha - 1)$ and with regularity condition $\alpha \notin \bigcup_{n \geq 0} E_n$, where $E_0 = \{\alpha \in \mathbb{C} : F(\alpha) = 0\}$, $F(\alpha) = \int_{-\infty}^{+\infty} e^{i\frac{\alpha^3}{3} - x^2 + i(\alpha-1)x} dx$, and for each integer $n \geq 1$, $E_n = \{\alpha \in \mathbb{C} : \Xi_n(\alpha) = 0\}$. Here, $\Xi_n(\alpha)$ is the Hankel determinant associated with u . Notice that u is a semi-classical linear functional of class one [10].

The recurrence coefficients β_n and γ_{n+1} , $n \geq 0$, of the sequence $\{B_n\}_{n \geq 0}$ are determined by the system [10] :

$$\begin{cases} \frac{n+1}{\gamma_{n+1}} &= 2 - i(\beta_n + \beta_{n+1}), \quad n \geq 0, \\ i(\gamma_{n+2} + \gamma_{n+1}) &= \psi(\beta_{n+1}), \quad n \geq 0, \\ \gamma_1 = -i\psi(\beta_0), \quad \beta_0 &= -i\frac{F'(\alpha)}{F(\alpha)}. \end{cases} \quad (4.10)$$

The sequence $\{B_n\}_{n \geq 0}$ is characterized by the following structure relation [10] :

$$B_n^{[1]}(x) = B_n(x) - \frac{i\gamma_n\gamma_{n+1}}{n+1}B_{n-1}(x), \quad n \geq 1. \quad (4.11)$$

Thus, taking into account $\lambda_{n,n-1} = -\frac{i\gamma_n\gamma_{n+1}}{n+1} \neq 0$, $n \geq 1$, we deduce a strictly finite-type relation between the sequences $\{B_n\}_{n \geq 0}$ and $\{B_n^{[1]}\}_{n \geq 0}$ with index $s = 1$, with respect to $\phi(x) = 1$,

From Proposition 4.1. we get the following inverse relation, for $n \geq 1$,

$$(x + v_{n,0}^*)B_n(x) = B_{n+1}^{[1]}(x) + \frac{i(n+1)}{\gamma_{n+1}}B_n^{[1]}(x), \quad (4.12)$$

where $v_{n,0}^* = \frac{i(n+1)}{\gamma_{n+1}} - \frac{i\gamma_{n+1}\gamma_{n+2}}{n+2} - \beta_n$, $n \geq 1$. The sequence $\{B_n\}_{n \geq 0}$ could be characterized by a relation as (4.12). It is the aim of the following result.

Proposition 4.2. *Let $\{B_n\}_{n \geq 0}$ be a MOPS satisfying (1.5). Then the following statements are equivalent.*

- i) There exists a set of non-zero complex numbers $\{\lambda_{n,n-1}\}_{n \geq 1}$ such that, for $n \geq 1$,
- $$B_n^{[1]}(x) = B_n(x) + \lambda_{n,n-1}B_{n-1}(x). \quad (4.13)$$

ii) There exists a set of complex numbers $\{\varrho_n\}_{n \geq 0}$, with $\varrho_n \neq 0$, $n \geq 1$, and $\varrho_0 = 0$, such that for $n \geq 0$,

$$(x + \frac{\gamma_{n+1}}{\varrho_{n+1}} + \varrho_n - \beta_n)B_n(x) = B_{n+1}^{[1]}(x) + \varrho_n B_n^{[1]}(x). \quad (4.14)$$

Proof. Assume that i) holds. From Proposition 4.1, we get

$$(x + \frac{\gamma_{n+1}}{\varrho_{n+1}} + \varrho_n - \beta_n)B_n(x) = B_{n+1}^{[1]}(x) + \varrho_n B_n^{[1]}(x), \quad n \geq 1,$$

where $\varrho_n = \gamma_n \lambda_{n,n-1}^{-1}$, $n \geq 1$. For $n = 1$, in (4.13), we obtain $\lambda_{1,0} = \frac{\beta_0 - \beta_1}{2}$. Then, $\frac{\gamma_1}{\varrho_1} = \frac{\beta_0 - \beta_1}{2}$. Hence,

$$(x + \frac{\gamma_1}{\varrho_1} - \beta_0)B_0(x) = x - \frac{\beta_0 + \beta_1}{2} = B_1^{[1]}(x) + \varrho_0 B_0^{[1]}(x),$$

i.e. $\varrho_0 = 0$. Thus, ii) holds. Conversely, let us take $\lambda_{n,n-1} = \frac{\gamma_n}{\varrho_n}$, $n \geq 1$, and consider the MPS $\{A_n\}_{n \geq 0}$ defined by

$$A_n(x) = B_n(x) + \lambda_{n,n-1}B_{n-1}(x), \quad n \geq 1. \quad (4.15)$$

From Proposition 4.1, we get

$$(x + v_{n,0}^*)B_n(x) = A_{n+1}(x) + \theta_{n,n}^*A_n(x), \quad n \geq 1,$$

where $v_{n,0}^* = \frac{\gamma_{n+1}}{\varrho_{n+1}} + \varrho_n - \beta_n$, $n \geq 1$, and $\theta_{n,n}^* = \frac{\gamma_n}{\lambda_{n,n-1}} = \varrho_n$, $n \geq 1$. From the assumption ii) and the previous relation, it follows that

$$A_{n+1}(x) + \varrho_n A_n(x) = B_{n+1}^{[1]}(x) + \varrho_n B_n^{[1]}(x), \quad n \geq 1.$$

Equivalently,

$$A_n(x) - B_n^{[1]}(x) = (\prod_{\nu=1}^n \varrho_\nu)(A_1(x) - B_1^{[1]}(x)) = 0, \quad n \geq 1.$$

But, from (4.15) for $n = 1$ we get $A_1(x) = x - \beta_0 + \frac{\gamma_1}{\varrho_1}$.

From (4.14), with $n = 0$, we get $B_1^{[1]}(x) = x - \beta_0 + \frac{\gamma_1}{\varrho_1}$.

Hence, $A_n(x) = B_n^{[1]}(x)$, $n \geq 0$. Thus according to (4.15), i) holds. \square

5. The case $(t, s) = (0, 2)$

Let $\{B_n\}_{n \geq 0}$ be a MOPS with respect to the linear functional u and satisfying (1.5). Consider the following finite-type relation between $\{B_n\}_{n \geq 0}$ and $\{Q_n\}_{n \geq 0}$, with index $s = 2$, with respect to $\phi(x) = 1$, for $n \geq 0$,

$$Q_n(x) = B_n(x) + \lambda_{n,n-1}B_{n-1}(x) + \lambda_{n,n-2}B_{n-2}(x), \quad (5.1)$$

$$\exists r \geq 2, \quad \lambda_{r,r-2} \neq 0. \quad (5.2)$$

From Lemma 2.1, for every system of complex numbers $(\theta_{n,\nu})_{\nu=n}^{n+2}$, $n \geq 0$, where $\theta_{n,n+2} = 1$, $n \geq 0$ and $\theta_{r,r} \neq 0$, there exists a unique MPS $\{\Omega_2(x; n)\}_{n \geq 0}$, where $\Omega_2(x; n) = x^2 + v_{n,1}x + v_{n,0}$, $n \geq 0$, and a unique system of complex numbers, $(\zeta_{n,\nu}^{[0]})_{\nu=n-2}^{n-1}$, $n \geq 0$, such that

$$\sum_{\nu=n}^{n+2} \theta_{n,\nu} Q_\nu(x) = \Omega_2(x; n)B_n(x) + \zeta_{n,n-1}^{[0]}B_{n-1}(x) + \zeta_{n,n-2}^{[0]}B_{n-2}(x), \quad n \geq 0, \quad (5.3)$$

where

$$\left\{ \begin{array}{l} \lambda_{n+2,n+1} + \theta_{n,n+1} = \beta_{n+1} + \beta_n + v_{n,1}, \quad n \geq 0, \\ \lambda_{2,0} + \theta_{0,1}\lambda_{1,0} + \theta_{0,0} = \gamma_1 + \beta_0(\beta_0 + v_{0,1}) + v_{0,0}, \\ \lambda_{n+2,n} + \theta_{n,n+1}\lambda_{n+1,n} + \theta_{n,n} = \gamma_{n+1} + \gamma_n + \beta_n(\beta_n + v_{n,1}) + v_{n,0}, \quad n \geq 1, \\ \theta_{n,n+1}\lambda_{n+1,n-1} + \theta_{n,n}\lambda_{n,n-1} = \gamma_n(\beta_n + \beta_{n-1} + v_{n,1}) + \zeta_{n,n-1}^{[0]}, \quad n \geq 1, \\ \theta_{n,n}\lambda_{n,n-2} = \gamma_n\gamma_{n-1} + \zeta_{n,n-2}^{[0]}, \quad n \geq 2, \end{array} \right. \quad (5.4)$$

The determinants associated with (5.1) – (5.2) are

$$\begin{aligned} \Delta_0(0, 2) &= \Delta_1(0, 2) = 0, \\ \Delta_n(0, 2) &= \lambda_{n,n-2}(\lambda_{n+1,n-1} - \gamma_n), \quad n \geq 2. \end{aligned} \quad (5.5)$$

Proposition 5.1. Let $\{B_n\}_{n \geq 0}$ be a MOPS and $\{Q_n\}_{n \geq 0}$ be the MPS satisfying (5.1) – (5.2). For every fixed integer $p \geq 2$, the following statements are equivalent

$$i) \lambda_{n,n-2}(\lambda_{n+1,n-1} - \gamma_n) \neq 0, \quad n \geq p.$$

As a consequence of Theorem 3.2, where $t = 0$ and $s = 2$, we have the following result

- ii) There exist a unique SCN $(\theta_{n,\nu}^*)_{\nu=n}^{n+2}$, $n \geq p$, with $\theta_{n,n+2}^* = 1$, $n \geq p$, and $\theta_{r,r}^* \neq 0$, if $p \leq r$, and there exists a unique MPS $\{\Omega_2^*(x; n)\}_{n \geq p}$, where $\deg \Omega_2^*(x; n) = 2$, $n \geq p$, such that, for $n \geq p$,

$$\Omega_2^*(x; n)B_n(x) = Q_{n+2}(x) + \theta_{n,n+1}^*Q_{n+1}(x) + \theta_{n,n}^*Q_n(x). \quad (5.6)$$

We write

$$\theta_{n,n+1}^* = \frac{[\lambda_{n,n-2}(\beta_{n-1} - \beta_{n+1} + \lambda_{n+2,n+1}) - \lambda_{n,n-1}\gamma_{n-1}]\gamma_n}{\lambda_{n,n-2}(\lambda_{n+1,n-1} - \gamma_n)},$$

$$\theta_{n,n}^* = \frac{\gamma_n\gamma_{n-1}}{\lambda_{n,n-2}},$$

$$\Omega_2^*(x; n) = x^2 + v_{n,1}^*x + v_{n,0}^*, \quad n \geq p, \quad (5.7)$$

where

$$v_{n,0}^* = \theta_{n,n}^* + (\lambda_{n+1,n} - \beta_n)\theta_{n,n+1}^* - \gamma_{n+1} - \gamma_n + \lambda_{n+2,n} + \beta_n(\beta_{n+1} - \lambda_{n+2,n+1}),$$

$$v_{n,1}^* = \theta_{n,n+1}^* - \beta_{n+1} - \beta_n + \lambda_{n+2,n+1}.$$

Example. Let $\{B_n\}_{n \geq 0}$ be the sequence of monic polynomials, orthogonal with respect to the linear functional u such that

$$\langle u, p \rangle = \int_{-\infty}^{+\infty} p(x)e^{-x^4} dx.$$

This sequence of polynomials was introduced by P. Nevai (see [15]) in the framework of the so-called Freud measures. These polynomials satisfy the three-term recurrence relation (1.5), with coefficients $\beta_n = 0$, $n \geq 0$, and where γ_{n+1} , $n \geq 0$, are given by a non-linear recurrence relation (see [3] and [15])

$$n = 4\gamma_n(\gamma_{n+1} + \gamma_n + \gamma_{n-1}), \quad n \geq 1,$$

with $\gamma_0 = 0$ and $\gamma_1 = \Gamma(3/4)\Gamma(1/4)$.

The sequence $\{B_n\}_{n \geq 0}$ satisfies the following structure relation (see [3])

$$B_n^{[1]}(x) = B_n(x) + \lambda_{n,n-2}B_{n-2}(x), \quad n \geq 2, \quad (5.8)$$

where

$$\lambda_{n,n-2} = \frac{4}{n+1}\gamma_{n+1}\gamma_n\gamma_{n-1} \neq 0, \quad n \geq 2.$$

From (5.3), with $Q_n(x) = B_n^{[1]}(x)$, $n \geq 0$, and the fact that the polynomial sequences $\{B_n\}_{n \geq 0}$ and $\{B_n^{[1]}\}_{n \geq 0}$

are symmetric, i.e., $B_n(-x) = (-1)^n B_n(x)$, $n \geq 0$, we get, for $n \geq 0$,

$$B_{n+2}^{[1]}(x) + \theta_{n,n}B_n^{[1]}(x) = (x^2 + v_{n,0})B_n(x) + \zeta_{n,n-2}^{[0]}B_{n-2}(x), \quad (5.9)$$

where

$$\begin{cases} \lambda_{2,0} + \theta_{0,0} & = \gamma_1 + v_{0,0}, \\ \lambda_{n+2,n} + \theta_{n,n} & = \gamma_{n+1} + \gamma_n + v_{n,0}, \quad n \geq 1, \\ \theta_{n,n}\lambda_{n,n-2} & = \gamma_n\gamma_{n-1} + \zeta_{n,n-2}^{[0]}, \quad n \geq 2. \end{cases} \quad (5.10)$$

Since we have $\lambda_{n,n-2}$, $n \geq 2$, the choice $\zeta_{n,n-2}^{[0]} = 0$, $n \geq 2$, is possible and yields the inverse relation

$$(x^2 + v_{n,0}^*)B_n(x) = B_{n+2}^{[1]}(x) + \theta_{n,n}^*B_n^{[1]}(x), \quad n \geq 0, \quad (5.11)$$

where

$$\theta_{n,n}^* = \frac{n+1}{4\gamma_{n+1}},$$

$$v_{n,0}^* = \frac{n+1}{4\gamma_{n+1}} - \gamma_n - \gamma_{n+1} + \frac{4}{n+3}\gamma_{n+1}\gamma_{n+2}\gamma_{n+3}.$$

Here, the determinants associated with (5.8) are

$$\Delta_n(0, 2) = \frac{4}{n+1}\gamma_{n+1}\gamma_n^2\gamma_{n-1} \left[\frac{4}{n+2}\gamma_{n+2}\gamma_{n+1} - 1 \right], \quad (5.12)$$

$n \geq 2$, with $\Delta_0(0, 2) = \Delta_1(0, 2) = 0$.

From Proposition 5.1, we deduce that the uniqueness of the previous inverse relation requires that $\lambda_{n+1,n-1} - \gamma_n = \gamma_n \left[\frac{4}{n+2}\gamma_{n+2}\gamma_{n+1} - 1 \right] \neq 0$, $n \geq 2$. Equivalently, $4\gamma_{n+2}\gamma_{n+1} \neq n+2$, $n \geq 2$. Indeed, by using (5.8), where n is replaced by $n+1$ and taking into account the orthogonality of the polynomial sequence $\{B_n\}_{n \geq 0}$, we get $B_{n+1}^{[1]}(x) = xB_n(x) + (\lambda_{n+1,n-1} - \gamma_n)B_{n-1}(x)$, $n \geq 1$. On the other hand, if we suppose that there exists an integer $n_0 \geq 2$ such that $\lambda_{n_0+1,n_0-1} - \gamma_{n_0} = 0$, then $B_{n_0+1}^{[1]}(x) = xB_{n_0}(x)$. In this case (5.11), with $n = n_0$ will be written as $(x^2 + \alpha x + v_{n,0}^*)B_n(x) = B_{n+2}^{[1]}(x) + \alpha B_{n+1}^{[1]}(x) + \theta_{n,n}^*B_n^{[1]}(x)$, for all $\alpha \in \mathbb{C}$. This contradicts the uniqueness of the inverse relation.

Acknowledgements: The first author (FM) was supported by Dirección General de Investigación (Ministerio de Educación y Ciencia) of Spain under grant MTM 2006-13000-C03-02. The work of second author (RS) was supported by Entreprise Kilani at Gabès and Institut Supérieur de Gestion de Gabès, Tunisie.

References

- [1] M. Alfaro, F. Marcellán, A. Peña & M. L. Rezola, *On linearly related orthogonal polynomials and their functionals*, J. Math. Anal. Appl. **287** (2003), 307–319.
- [2] M. Alfaro, F. Marcellán, A. Peña & M. L. Rezola, *On rational transformations of linear functionals. A direct problem*, J. Math. Anal. Appl. **298** (2004), 171–183.
- [3] A. Cachafeiro, F. Marcellán & J. J. Moreno-Balcázar, *On asymptotic properties of Freud-Sobolev orthogonal polynomials*, J. Approx. Theory **125** (2003), 26–41.
- [4] T. S. Chihara, "An Introduction to Orthogonal Polynomials", Gordon and Breach, New York, 1978.
- [5] A. M. Delgado & F. Marcellán. *Companion linear functionals and Sobolev inner products: a case study*, Meth. Appl. Anal. **11** (2004), 237–266.
- [6] F. Marcellán, M. Alfaro & M. L. Rezola, *Orthogonal polynomials on Sobolev spaces: Old and new directions*, J. Comput. Appl. Math. **48** (1993), 113–131.
- [7] F. Marcellán & J. C. Petronilho, *Orthogonal polynomials and coherent pairs: The classical case*, Indag. Math. (NS), **6** (1995), 287–307.
- [8] F. Marcellán, T. E. Pérez & M. A. Piñar, *Orthogonal polynomials on weighted Sobolev spaces: the semiclassical case*, Ann. Numer. Math. **2** (1995), 93–122.
- [9] F. Marcellán & J. C. Medem, *Q-Classical orthogonal polynomials: a very classical approach*, Elect. Trans. on Numer. Anal. **9** (1999), 112–127.
- [10] P. Maroni, *Un exemple d'une suite orthogonal semi-classique de classe un*. In Publ. Labo. d'Analyse Numérique, Université Pierre et Marie Curie, Paris. **89033** (1989).
- [11] P. Maroni, *Fonctions eulériennes. Polynômes orthogonaux classiques*. In Techniques de l'ingénieur, A **154** (1994), 1–30.
- [12] P. Maroni, *Semi-classical character and finite-type relations between polynomial sequences*. J. Appl. Num. Math. **31** (1999), 295–330.
- [13] P. Maroni & R. Sfaxi, *Diagonal orthogonal polynomial sequences*, Meth. Appl. Anal. **7** (2000), 769–792.
- [14] H. G. Meijer, *Determination of all Coherent Pairs*. J. Approx. Theory, **89** (1997), 321–343.
- [15] P. Nevai, *Orthogonal Polynomials associated with $\exp(-x^4)$* . Proc. Canad. Math. Soc. **3** (1983), 263–285.

Recibido el 16 de diciembre de 2007

Aceptado para su publicación el 14 de marzo de 2008